



LKB1 is physiologically required for sleep from *Drosophila melanogaster* to the *Mus musculus*

Ziyi Liu ^{1,2,3,4} Lifan Jiang,⁵ Chaoyi Li,⁵ Chengang Li,^{1,2,3,4} Jingqun Yang,^{1,2,3,4} Jianjun Yu,^{1,2,3,4} Renbo Mao,^{1,2,3,4} Yi Rao ^{1,2,3,4,*}

¹Peking University-Tsinghua University-National Institute of Biological Sciences Joint Graduate Program, School of Life Sciences, PKU-IDG/McGovern Institute for Brain Research, School of Chemistry and Molecular Engineering, School of Pharmaceutical Sciences, Peking University, Beijing 100871, China

²Chinese Institute for Brain Research, Beijing, China

³Capital Medical University, Beijing, China

⁴Changping Laboratory, Beijing, China

⁵Shenzhen Bay Laboratory, Institute of Molecular Physiology, Shenzhen, Guangdong, China

*Corresponding author: Peking University-Tsinghua University-National Institute of Biological Sciences Joint Graduate Program, School of Life Sciences, PKU-IDG/McGovern Institute for Brain Research, School of Chemistry and Molecular Engineering, School of Pharmaceutical Sciences, Peking University, Beijing 100871, China. Email: yrao@pku.edu.cn

Abstract

LKB1 is known as a master kinase for 14 kinases related to the adenosine monophosphate (AMP)-activated protein kinase (AMPK). Two of them (SIK3 and AMPK α) have previously been implicated in sleep regulation. We generated loss-of-function (LOF) mutants for Lkb1 in both *Drosophila* and mice. Sleep, but not circadian rhythms, was reduced in Lkb1-mutant flies and in flies with neuronal deletion of Lkb1. Genetic interactions between Lkb1 and AMPK T184A mutants in *Drosophila* sleep or those between Lkb1 and SIK3 T196A mutants in *Drosophila* viability have been observed. Sleep was reduced in mice after virally mediated reduction of Lkb1 in the brain. Electroencephalography (EEG) analysis showed that non-rapid eye movement (NREM) sleep and sleep need were both reduced in Lkb1-mutant mice. These results indicate that LKB1 plays a physiological role in sleep regulation conserved from flies to mice.

Keywords: sleep; LKB1; flies; mice

Introduction

Human Peutz-Jeghers syndrome (PJS) (Peutz 1921; Jeghers et al. 1949) is an autosomal dominant disorder with gastrointestinal (GI) polyps and increased cancer risk of multiple tissues (Tomlinson and Houlston 1997; Westerman et al. 1999). The gene mutated in, and responsible for, PJS encodes the liver kinase B1 (LKB1, also known as STK11) (Hemminki et al. 1997; 1998; Jenne et al. 1998). Lkb1 is thus a tumor suppressor gene, mutated in multiple cancers, especially the GI (Mehenni et al. 1998; Bardeesy et al. 2002; Jishage et al. 2002; Miyoshi et al. 2002; Hearle et al. 2006) and lung adenocarcinoma (Sanchez-Cespedes et al. 2002; Carretero et al. 2004; Ji et al. 2007; Matsumoto et al. 2007; Gill et al. 2011; Skoulidis et al. 2015), cervical cancer (Wingo et al. 2009), ovarian cancer (Tanwar et al. 2014), breast cancer (Shen et al. 2002; Hearle et al. 2006; Sengupta et al. 2017), pancreatic cancer (Morton et al. 2010), and melanoma (Guldberg et al. 1999; Rowan et al. 1999).

LKB1 phosphorylates threonine 172 (T172) of the α subunit of adenosine monophosphate (AMP)-activated protein kinase (AMPK α) (Hawley et al. 2003; Hong et al. 2003; Sutherland et al. 2003; Woods et al. 2003; Lizcano et al. 2004; Shaw et al. 2004, 2005; Sakamoto et al. 2005), and positively regulates the activity of AMPK.

AMPK is a well-known kinase (Beg et al. 1973; Carlson and Kim 1973; Ingebritsen et al. 1978; Yeh and Kim 1980; Ferrer et al. 1985; Carling et al. 1987, 1989; Munday et al. 1988) with important physiological and pathological roles (Hardie 2014; Lopez and Dieguez 2014; Hardie et al. 2016; Herzig and Shaw 2018). The α , β , and γ subunits of AMPK form a heterotrimeric complex (Davies et al. 1994; Mitchelhill et al. 1994; Michell et al. 1996). The catalytic α subunit is regulated by phosphorylation at T172 of AMPK α 2 or T183 of AMPK α 1 (Hawley et al. 1996).

There are 12 additional mammalian AMPK-related kinases (ARKs), similar to the α subunit of AMPK, all regulated at the site equivalent to AMPK-T172 (Lizcano et al. 2004). LKB1 and its associated proteins STE20-related adaptor (STRAD) and mouse protein 25 (MO25) have been reported to phosphorylate all 14 ARKs (Lizcano et al. 2004), making LKB1 a master kinase for ARKs (Lizcano et al. 2004; Alessi et al. 2006; Shackelford and Shaw 2009). We have recently found that more than 20 kinases in the STE20 family of mammalian serine-threonine kinases could phosphorylate ARKs in vitro, though the physiological roles of STE20 kinases in ARK phosphorylation remain unknown (Liu, Wang, Cui, Gao, et al. 2022; Liu, Wang, Cui, et al. 2022).

Received: December 13, 2021. **Accepted:** May 10, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of Genetics Society of America. All rights reserved.

For permissions, please email: journals.permissions@oup.com

Some ARKs have been reported to regulate sleep. In mice, inhibitors of AMPK were found to decrease sleep, whereas activators of AMPK were found to increase sleep (Chikahisa et al. 2009). In flies, knockdown of AMPK β in neurons decreased the total amount of sleep and resulted in fragmented sleep (Nagy et al. 2018). Knockdown of AMPK α in a specific pair of neurons suppressed sleep (Yurgel et al. 2019).

Studies in mice have shown that sleep was increased in gain-of-function (GOF) mutations in the salt inducible kinase (SIK) 3 (Funato et al. 2016), and sleep need was reduced in GOF mutants of SIK 1, 2, and 3 (Funato et al. 2016; Honda et al. 2018; Park et al. 2020). Sleep was also decreased when SIK3 was downregulated in flies (Funato et al. 2016). Here we investigated the functional role of LKB1 in regulating sleep in flies and mice.

Materials and methods

Fly lines and rearing conditions

Flies were reared on standard corn meal at 25°C, 60% humidity and kept in 12 hours (h) light/12 h dark (LD) conditions. 57C10-Gal4, nos- ϕ C31, hs-Cre on X were from the Bloomington Stock Center. vas-Cas9 was a gift from Dr J. Ni (Tsinghua University, Beijing). Upstream activation sequence (UAS)-Cas9, UAS-Ampk, UAS-Ampk-T184A, UAS-Ampk-T184E, Sik3-flag, Sik3-T196A-flag, and Sik3-T196E-flag flies were from our laboratory.

Flies used in behavioral assays were backcrossed into our isogenized Canton S background for 7 generations. All results of sleep analysis in this paper were obtained from female flies.

Generation of KO, KI, and transgenic lines

Total RNA was extracted from isoCS by TRIzol reagent (Invitrogen). Using the PrimeScript II 1st Strand cDNA Synthesis Kit (Takara), we reverse-transcribed the extracted mRNA into cDNA. The UAS-Lkb1 flies was constructed by inserting the coding sequence (CDS) of CG9374 amplified from cDNA into the pACU2 plasmid (a gift from the Jan Lab at UCSF) (Han et al. 2011) before being inserted into the attP40 site.

The UAS-Lkb1-sgRNA construct was designed by inserting the sgRNAs into pMt: sgRNA^{3XEF} vectors based on pACU2, with rice tRNA separating the different sgRNAs. CRISPR-Gold website was used to design 3 sgRNAs of Lkb1 (Supplementary Fig. 3) (Chu et al. 2016; Poe et al. 2019). The construct was inserted into the attP40 site.

KO and KI lines were generated as described previously (Deng et al. 2019). Knockout flies were generated with the CRISPR/Cas9 system. Two different sgRNAs were constructed with U6b-sgRNA plasmids. The 5' homologous arm and the 3' homologous arm of ~2 kb amplified from the wt fly genome were inserted into a pBSK plasmid for homologous recombination repair. The cassette of attP-3P3-RFP was introduced in the middle. sgRNA plasmids and the donor plasmids were injected into vas-Cas9 embryos to introduce attP-3P3-RFP into the genome at the region of interest and replaced it by homologous recombination. 3P3-RFP served as a marker to screen for the correct flies. Primers across the homologous arms were designed to verify the sequences by PCR and DNA sequencing. attP site was introduced into the genome with 3P3-RFP-LoxP. For KI files, the nos- ϕ C31 virgin females were first crossed with knock-out males and the pBSK plasmid inserted with attB-T2A-Gal4-miniwhite-LoxP cassette was injected into the female embryos. Miniwhite serves as a marker to screen for the correct flies, which could be excised by the Cre/LoxP system. Primers were designed to verify the sequence by PCR and DNA sequencing.

Quantitative PCR

Total RNA was extracted from 30 flies of 5–7 days old by TRIzol reagent (Invitrogen). The genomic template was removed using DNase (Takara). cDNA was reverse-transcribed using Takara's PrimeScript II 1st Strand cDNA synthesis kit (Takara). Quantitative PCR was carried out with TransStart Top Green qPCR SuperMix kit (TransGen) in the Bio-Rad PCR system (CFX96 Touch Deep Well). The sequences of primers to detect Lkb1 and Rp49(endogenous control) mRNA are

Lkb1-F: 5'-GCCGTCAAGATCCTGACTA-3'
Lkb1-R: 5'-CTCCGCTGGACCAGATG-3'
Rp49-F: 5'-CGACGCTTCAAGGGACAGTATC-3'
Rp49-R: 5'-TCCGACCAGGTTACAAGAACTCTC-3'

Drosophila sleep analysis

Drosophila sleep analysis was performed as described previously (Qian et al. 2017; Dai et al. 2019). 5–7 days old flies were placed in a 65 mm \times 5 mm clear glass tube with one end containing food and the other end plugging with cotton. All flies were recorded by video-cameras. Before sleep measurement, flies were entrained to an LD cycle at 25°C, 60% humidity for at least 2 days, and infrared LED light was used to ensure constant illumination when lights off. Immobility longer than 5 min was defined as one sleep event (Hendricks et al. 2000; Shaw et al. 2000). Information of fly location was tracked and sleep parameters were analyzed using Matlab (Mathworks), from which dead flies were removed. Sleep duration, sleep bout duration, sleep bout number, and sleep latency for each LD were analyzed. Each experiment was repeated at least 3 times.

Drosophila circadian analysis

Flies were reared and recorded in the same condition as sleep assay as described in papers from our laboratory (Qian et al. 2017; Dai et al. 2021), except that the condition was constant darkness. Six to 8 days activity was measured and calculated in ActogramJ (Klarsfeld et al. 2003). Rhythmic strength, power and period were calculated by Chi-square method.

Immunoblot analysis

Mouse brains were quickly dissected and washed with phosphate buffer saline on ice. Lysis buffer (20 mM HEPES, 10 mM KCl, 1.5 mM MgCl₂, 1 mM EDTA, 1 mM EGTA, 1 mM DTT, freshly supplemented with a protease and phosphatase inhibitors cocktail) were used to homogenize brains by homogenizer (Wiggins, D-500 pro) at 4°C. Brain homogenates were centrifuged at 14,000 revolution per minutes for 15 min at 4°C. The supernatant was transferred to a new microtube and quantified with bicinchoninic acid assay (Thermo Fisher, 23225). The supernatant was analyzed by SDS-PAGE and proteins were transferred to a nitrocellulose membrane (GE Healthcare, #BA85). Membranes were incubated for 1 h in a blocking solution (tris-buffered saline containing 0.1% Tween-20, 5% milk). Primary antibodies were anti-LKB1 (cell signaling, #3047) and anti-ACTIN (Santa Cruz, sc-8342).

Retro-orbital injection in mice

Mice were reared at controlled temperature and humidity conditions with 12 h light/12 h dark cycle. Food and water were provided ad libitum. Lkb1^{fl/fl} mice were from the Jackson Laboratory (JAX #014143). They contained loxP sites flanking exons 3–6 of Lkb1 gene (Nakada et al. 2010). adeno-associated viral (AAV)-PHP.eB-hSyn-Cre-green fluorescent protein (GFP) and

AAV-PHP.eB-hSyn-GFP virus were from Chinese Institute for Brain Research, Beijing. All results of sleep analysis in this paper were obtained from female mice.

Total 0.2 ml/10 g avertin was injected intraperitoneally into the mice for anesthetization. Rodent eyes were protruded by gentle downward pressure to the skin on the dorsal and ventral sides of the eye. The operator inserted the needle beveled downward into the retro-orbital sinus at the medial corner of the eye (Yardeni et al. 2011). The AAV-PHP.eB virus was injected for whole brain infection (Chan et al. 2017).

Mouse sleep analysis

Mouse sleep analysis was described in a previous article from our laboratory (Zhang et al. 2018). Eight-week-old mice were selected for retro-orbital injection. One week after viral injection, EEG and EMG electrode implantation procedures were performed. Mice were allowed to recover for more than 5 days individually and placed in a recording cage and tethered to an omni-directional arm (RWD Life Science Inc.) with connection cable for 2 days of habituation before recording. EEG and EMG data were recorded with custom software at a sampling frequency of 200 Hz for 2 consecutive days to analyze sleep/wake behavior under baseline conditions. The recording chamber was maintained at 12 h LD cycle and controlled temperature (24–25°C). EEG/EMG data were initially processed by Accusleep (Barger et al. 2019) before manual correction in SleepSign to improve accuracy. WAKE was scored as high amplitude and variable EMG and fast and low amplitude EEG. Nonrapid eye movement (NREM) was scored as high amplitude δ (1–4 Hz) frequency EEG and low EMG tonus. REM was scored as a complete silent of EMG signals and low amplitude high frequency θ (6–9 Hz)-dominated EEG signal.

For power spectrum analysis, EEG was subjected to fast Fourier transform analysis with a Hamming window method by SleepSign, yielding power spectra between 0 and 25 Hz with a 0.39 Hz bin resolution. Epochs containing movement artifacts were marked, included in sleep duration analysis but excluded from the power spectra analysis. Power spectra for each vigilance state represents the mean power distribution of this state during a 24-h baseline recording. The δ -power density of NREMs per hour represents the average of δ -power density as a percentage of δ -band power (1–4 Hz) to total power (0–25 Hz) for each NREM epoch contained in an hour.

Statistics

All statistical analyses were performed with Prism 7 (GraphPad Software). Differences in means between samples larger than 2 groups were analyzed using ordinary 1-way ANOVA. Unpaired t test was used for 2 groups comparison. Power spectrum between different lines was compared by 2-way ANOVA followed by Turkey's multiple comparisons test. n.s. denotes $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; and *** $P < 0.001$ for all statistical results in this paper.

Results

Sleep phenotypes of *Drosophila* Lkb1 mutants

Null mutants for Lkb1 are lethal in *Drosophila* (Martin and St Johnston 2003). We had generated a Lkb1 knockout ("lkb1^{T1}") line (Supplementary Figs. 1a and 3b) and found that lkb1^{T1/T1} mutation was lethal in the pupa stage. The level of Lkb1 mRNA was reduced in the heterozygous lkb1^{T1/+} flies (Supplementary Fig. 1b). We then tested whether the heterozygous lkb1^{T1} had any phenotype in sleep using flies kept in 12 hours (h) light/12 h dark (LD)

cycles (Supplementary Fig. 1c). While lkb1^{T1/+} flies were not significantly different from the wild-type (wt) flies in sleep bout numbers (Supplementary Fig. 1e), or daytime sleep duration (Supplementary Fig. 1d), daytime sleep bout duration (Supplementary Fig. 1f), lkb1^{T1/+} flies showed significantly lower nighttime sleep duration (Supplementary Fig. 1d), nighttime sleep bout duration (Supplementary Fig. 1f), and longer latency to sleep (Supplementary Fig. 1g). Thus, there was dosage-sensitive physiological requirement of Lkb1 in nighttime sleep.

We tried to, and succeeded in, generating lkb1^{T2}, a hypomorphic mutation for Lkb1 in flies (Fig. 1a; Supplementary Fig. 3, a and b). Lkb1 mRNA was significantly reduced in lkb1^{T2/+} and lkb1^{T2/T2} flies (Fig. 1b). During the day, lkb1^{T2/T2} flies were not significantly different from the lkb1^{T2/+} and wt flies in sleep duration (Fig. 1, c and d), sleep bout number (Fig. 1e), sleep bout duration (Fig. 1f), or latency to sleep (Fig. 1g). During the night, not only lkb1^{T2/T2} flies showed significantly reduced sleep duration (Fig. 1, c and d), highly reduced sleep bout duration (Fig. 1f) and highly increased latency (Fig. 1g) than the wt flies, but also the heterozygous lkb1^{T2/+} flies were significantly different from the wt flies in all these parameters (Fig. 1, c–g), indicating a dosage sensitive requirement for Lkb1.

We examined the phenotypes of lkb1^{T1/T2}. Consistent with the lkb1^{T2/T2}, the mRNA levels of Lkb1 were significantly reduced in lkb1^{T1/T2} compared with wt, lkb1^{T1/+} and lkb1^{T2/+}, and even lower than that in lkb1^{T2/T2} (Supplementary Fig. 2a).

The sleep phenotype in lkb1^{T1/T2} flies was also consistent with lkb1^{T2/T2}, with highly reduced nighttime sleep duration (Supplementary Fig. 2, b and c), highly reduced sleep bout duration (Supplementary Fig. 2e) and highly increased latency to sleep (Supplementary Fig. 2f), when compared with wt, lkb1^{T1/+}, and lkb1^{T2/+} flies.

Results of sleep analysis of lkb1^{T1/+}, lkb1^{T2/+}, lkb1^{T2/T2}, and lkb1^{T1/T2} mutant flies all consistently support that Lkb1 plays a physiological role in promoting sleep.

Rescue of sleep phenotypes by Lkb1 in flies

We inserted the sequence of the yeast transcription factor Gal4 into the lkb1^{T2} mutant flies, flanking the lkb1 promoter, and obtained lkb1^{T2}-Gal4 flies (Fig. 2a). We also generated UAS-Lkb1 flies in which the Lkb1 CDS was expressed under the control of the UAS (Brand and Perrimon 1993). Because Gal4 protein binds to the UAS, the expression of Lkb1 in flies resulting from the crosses between lkb1^{T2}-Gal4 flies and UAS-Lkb1 flies would be under the control of the endogenous Lkb1 promoter. Indeed, expression of Lkb1 mRNA was restored when lkb1^{T2}-Gal4 and UAS-Lkb1 were present in the same flies (Fig. 2b), whereas Lkb1 mRNA was less in UAS-Lkb1; lkb1^{T2/T2} mutant flies, and lkb1^{T2}-Gal4/lkb1^{T2}-Gal4 flies than that in the wt. UAS-Lkb1 alone could not restore Lkb1 mRNA expression level to that in wt flies (Fig. 2b).

Both daytime and nighttime sleep durations were less in lkb1^{T2}-Gal4/lkb1^{T2}-Gal4 flies than those in wt flies (Fig. 2c). Introduction of UAS-Lkb1 in lkb1^{T2/T2} flies or lkb1^{T2}-Gal4 alone could not restore sleep. When both lkb1^{T2}-Gal4 and UAS-Lkb1 were present, nighttime sleep durations were restored (Fig. 2d). Nighttime sleep bout number, nighttime sleep bout duration, and nighttime latency were restored when both lkb1^{T2}-Gal4 and UAS-Lkb1 were present, but not when lkb1^{T2}-Gal4 or UAS-Lkb1 alone was present (Fig. 2, e–g).

These results support that the sleep phenotypes of lkb1^{T2/T2} were attributable to the reduction of Lkb1 mRNA expression in these flies.




Fig. 1. Sleep phenotypes of *Lkb1* knock-down mutant flies. a) A diagram illustrating the *Lkb1* insertion mutant *lkb1*^{T2}. b) Relative *Lkb1* mRNA levels in *lkb1*^{T2/T2} (red), *lkb1*^{T2/+} (blue), and wt (black) flies. c) Sleep profiles of *lkb1*^{T2/T2} (red, *n* = 42), *lkb1*^{T2/+} (blue, *n* = 44), and wt (black, *n* = 44) flies in a 12 h light/12 h dark (LD) cycle. Statistical analysis of sleep duration, sleep bout number, sleep bout duration, and latency to sleep in *lkb1*^{T2/T2} (red, *n* = 42), *lkb1*^{T2/+} (blue, *n* = 44), and wt (black, *n* = 44) flies. Open bars denote daytime, filled bars denote nighttime. d) Sleep duration. Nighttime sleep durations of *lkb1*^{T2/T2} mutants were significantly less than those in *lkb1*^{T2/+} and wt flies. e) Sleep bout number. Daytime sleep bout number of *lkb1*^{T2/T2} mutants was less than that of wt flies. f) Sleep bout duration. Nighttime sleep bout duration of *lkb1*^{T2/T2} mutants was significantly less than those of *lkb1*^{T2/+} and wt flies. g) Latency to sleep. Latency to sleep after light-off of *lkb1*^{T2/T2} mutants was significantly prolonged than *lkb1*^{T2/+} and wt flies. One-way ANOVA was used. n.s. denotes *P* > 0.05, **P* < 0.05, ***P* < 0.01, ****P* < 0.001. Error bars represent standard error of the mean (SEM).

Sleep phenotypes of flies carrying neuronal deletion of the *Lkb1* gene

To determine whether *Lkb1* functions in neurons, we used the CRISPR-Cas9 system to delete *Lkb1* from neurons specifically (Supplementary Fig. 4). A pan-neuronal Gal4 driver (57C10-Gal4) was used to control the expression of small guide RNA (sgRNA) targeting *Lkb1* in neurons. Compared with 57C10-Gal4>UAS-Cas9 alone or 57C10-Gal4>UAS-*Lkb1*-sgRNA alone, when both UAS-Cas9 and UAS-*Lkb1*-sgRNA were present in flies, nighttime sleep duration (Fig. 3b) and nighttime sleep bout duration (Fig. 3d) were significantly reduced and nighttime sleep latency significantly lengthened (Fig. 3e). Daytime sleep duration, bout

number, bout duration and latency were not significantly affected by neuronal gene targeting of *Lkb1* (Fig. 3, b–e).

We also investigated any potential effect that overexpression of *Lkb1* in neurons might cause (Supplementary Fig. 5a). We detected no phenotype resulting from neuronal overexpression of *Lkb1* on daytime and nighttime sleep duration, sleep bout number, sleep bout duration, or latency (Supplementary Fig. 5, b–f).

In all 3 series of experiments (Figs. 1–3), nighttime sleep phenotypes were more obvious than daytime sleep phenotypes. These results strongly indicate that *Lkb1* expression in neurons are required physiologically for sleep, especially nighttime sleep.




Fig. 2. Rescue of sleep loss in *lkb1^{T2/T2}* by Lkb1. a) A diagram of *lkb1^{T2}-Gal4*: a cDNA for the yeast Gal4 gene inserted in the *lkb1^{T2}* knockdown mutants. b) Relative Lkb1 mRNA levels in *lkb1^{T2}-Gal4* (blue), UAS-Lkb1;*lkb1^{T2}-Gal4* (red), UAS-Lkb1;*lkb1^{T2}* (yellow), and wt (black) flies. In *lkb1^{T2}-Gal4* homozygous flies, UAS-Lkb1 cDNA driven by Gal4 to rescue sleep phenotypes of *lkb1* knockdown mutants. c) Sleep profiles of UAS-Lkb1;*lkb1^{T2}-Gal4* (red, $n = 45$), UAS-Lkb1;*lkb1^{T2}* (yellow, $n = 47$), *lkb1^{T2}-Gal4* (blue, $n = 46$), and wt (black, $n = 36$) flies. Statistical analysis of sleep duration, sleep bout number, sleep bout duration and latency to sleep in UAS-Lkb1;*lkb1^{T2}-Gal4* (red, $n = 45$), UAS-Lkb1;*lkb1^{T2}* (yellow, $n = 47$), *lkb1^{T2}-Gal4* (blue, $n = 46$), and wt (black, $n = 36$) flies. Open bars denote daytime, filled bars nighttime. d) Sleep duration. Nighttime sleep duration of UAS-Lkb1;*lkb1^{T2}-Gal4* was similar to that of wt mutants, both significantly higher than UAS-Lkb1;*lkb1^{T2}* and *lkb1^{T2/T2}-Gal4* flies. e) Sleep bout number. Nighttime sleep bout number of UAS-Lkb1;*lkb1^{T2}-Gal4* was similar to the wt but significantly higher than UAS-Lkb1;*lkb1^{T2}* and *lkb1^{T2}-Gal4* flies. f) Sleep bout duration. Nighttime sleep bout duration of UAS-Lkb1;*lkb1^{T2}-Gal4* was similar to the wt but significantly higher than UAS-Lkb1;*lkb1^{T2}* and *lkb1^{T2}-Gal4* flies. g) Latency to sleep. Latency after light-off of UAS-Lkb1;*lkb1^{T2}-Gal4* was similar to the wt but significantly shorter than UAS-Lkb1;*lkb1^{T2}* and *lkb1^{T2}-Gal4* flies. One-way ANOVA was used. n.s. denotes $P > 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Error bars represent SEM.

Genetic interactions between Lkb1 and Ampk or Sik3 in flies

To examine potential genetic interactions of Lkb1 with either Ampk or Sik3, we combined the loss-of-function (LOF) Lkb1 mutation *lkb1^{T2}* with specific point mutations in either Ampk or Sik3.

The regulatory site T184 in *Drosophila* AMPK and T196 in *Drosophila* SIK3 were equivalent to T172 of mammalian AMPK2 and T221 of mammalian SIK3, important for their activities. When the endogenous T184 in fly AMPK was mutated to alanine (A) or glutamic acid (E), flies were lethal. We therefore introduced T184A and T184E mutations into an Ampk transgene whose expression was controlled by UAS. We introduced UAS-Ampk, UAS-Ampk-T184A, and UAS-Ampk-T184E into *lkb1^{T2/T2}* flies and used a pan-neuronal driver to express them in neurons (Fig. 4). Neuronal overexpression of Ampk-T184E (Fig. 4a) and UAS-Ampk (Fig. 4c) in *lkb1^{T2}* flies did not significantly change the sleep phenotypes of *lkb1^{T2/T2}* flies, but neuronal overexpression of

UAS-Ampk-T184A (Fig. 4b) in *lkb1^{T2/T2}* flies further decreased nighttime sleep duration.

Point mutations of Sik3-flag, Sik3-T196A-flag, and Sik3-T196E-flag were constructed in *Drosophila*. When the endogenous T196 in Sik3 was mutated to A or E, we could get homozygous flies. Upon crossing to *lkb1^{T2/T2}*, Sik3-T196A-flag; *lkb1^{T2/T2}* were homozygous lethal. The cross of Sik3-T196E-flag into the *lkb1^{T2/T2}* background generated viable flies, with no detectable change in sleep (Fig. 5).

Allele-specific genetic interactions between Lkb1 and Ampk in sleep, or between Lkb1 and Sik3 in viability, suggest, but do not prove, regulatory relationships between Lkb1 and Ampk in sleep or Sik3 in viability.

Circadian rhythm in Lkb1 mutant flies

The transcription factor differentiated embryo-chondrocyte 1 regulates circadian rhythm and can negatively regulate the




Fig. 3. Sleep phenotypes of mutants from whose neurons Lkb1 was targeted. a) Sleep profiles of UAS-Lkb1-sgRNA/57C10-Gal4;+/UAS-Cas9 (red, n = 44), UAS-Lkb1-sgRNA/57C10-Gal4 (blue, n = 41), and 57C10-Gal4/+;+/UAS-Cas9 (black, n = 45) flies. Statistical analysis of sleep duration, sleep bout number, sleep bout duration, and latency to sleep in UAS-Lkb1-sgRNA/57C10-Gal4;+/UAS-Cas9 (red, n = 44), UAS-Lkb1-sgRNA/57C10-Gal4 (blue, n = 41), and 57C10-Gal4/+;+/UAS-Cas9 (black, n = 45) flies. Open bars denote daytime, filled bars nighttime. b) Sleep duration. Daytime and nighttime sleep duration of UAS-Lkb1-sgRNA/57C10-Gal4;+/UAS-Cas9 was significantly less than those of UAS-Lkb1-sgRNA/57C10-Gal4 and 57C10-Gal4/+;+/UAS-Cas9 flies. c) Sleep bout number. Daytime and nighttime sleep bout numbers of UAS-Lkb1-sgRNA/57C10-Gal4;+/UAS-Cas9 was not significantly different from those of UAS-Lkb1-sgRNA/57C10-Gal4 and 57C10-Gal4/+;+/UAS-Cas9 flies. d) Sleep bout duration. Nighttime sleep bout duration of UAS-Lkb1-sgRNA/57C10-Gal4;+/UAS-Cas9 was significantly less than that of UAS-Lkb1-sgRNA/57C10-Gal4 and 57C10-Gal4/+;+/UAS-Cas9 flies. e) Latency to sleep. Latency to sleep after light-off of UAS-Lkb1-sgRNA/57C10-Gal4;+/UAS-Cas9 was longer than that of 57C10-Gal4/+;+/UAS-Cas9 which was not significantly different from that of UAS-Lkb1-sgRNA/57C10-Gal4 flies. One-way ANOVA was used. n.s. denotes $P > 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Error bars represent SEM.

transcription of Lkb1 and subsequently reduce AMPK activity (Sato *et al.* 2015).

We tested whether the circadian rhythm was affected in Lkb1 mutant flies. Lkb1 mutant flies were not different from wt flies in period length (Supplementary Fig. 7b). Relative rhythmic power was increased in $lkb1^{T2/+}$ and $lkb1^{T2/T2}$ mutants than wt flies (Supplementary Fig. 7).

Sleep phenotypes in Lkb1 conditional knockout mice

To investigate potential involvement of Lkb1 in regulating sleep of mammalian animals, we obtained $Lkb1^{fl/fl}$ mice in which the loxP sites flanked exons 3–6 of the Lkb1 gene (Nakada *et al.* 2010). To delete the Lkb1 gene from these mice, we injected AAV constructs expressing either the Cre recombinase together with the GFP or GFP alone to infect the mouse brain. Cre-GFP or GFP was under the control of a neuronal specific promoter human Synapsin I (hSyn) (in AAV-PHP.eB-hSyn-Cre-GFP or AAV-PHP.eB-hSyn-GFP).

We analyzed the expression of LKB1 protein in mice (Fig. 6, a and b). Injection of Cre-GFP expressing virus into wt or $Lkb1^{fl/+}$ mice did not reduce LKB1 protein expression in the brain. Neither did injection of only GFP expressing virus into $Lkb1^{fl/fl}$ mice. This conclusion was reached by examination of either several mouse brains combined (Fig. 6a), or individual mouse brains (Fig. 6b).

Functionally, only when the Cre-GFP expressing virus was injected into $Lkb1^{fl/fl}$ mice, wake duration was significantly increased during daytime (Fig. 6c; Supplementary Fig. 6a), NREM sleep duration was significantly decreased during daytime (Fig. 6e; Supplementary Fig. 6b). Controls (Cre-GFP injection into wt or $Lkb1^{fl/+}$ mice, GFP injection into $Lkb1^{fl/fl}$ mice) did not significantly change any sleep phenotypes.

REM sleep duration was not significantly affected by Cre-GFP injection into $Lkb1^{fl/fl}$ mice (Fig. 6g; Supplementary Fig. 6c).

Power density in the 1–4 Hz range (δ power density) of NREM is a commonly accepted indicator of sleep need (Borbely *et al.* 1981; Borbely 1982; Daan *et al.* 1984; Tobler and Borbely 1986; Dijk *et al.* 1987; Werth *et al.* 1996; Franken *et al.* 2001). We found that NREM δ power density was significantly reduced when the Cre-GFP expressing virus was injected into $Lkb1^{fl/fl}$ mice (Fig. 6f). Analysis over 24 h indicated that significant reduction was observed over most of the daily cycle (Fig. 6i).

Discussion

Our results indicate that LKB1 is required for sleep regulation: it plays an important and conserved role by promoting sleep in both flies and mice. LKB1 plays this role in neurons in both species because gene targeting of Lkb1 in neurons led to reduction of sleep. In mice, with the additional advantage of EEG analysis, we




Fig. 4. Genetic interactions between *lkb1* and *ampk*. a) Sleep profiles of UAS-AMPK-T184E/57C10; *lkb1^{T2}* (red, $n = 19$), 57C10; *lkb1^{T2}* (yellow, $n = 24$), UAS-AMPK-T184E; *lkb1^{T2}* (blue, $n = 22$) and wt (black, $n = 24$) flies. b) Sleep profiles of UAS-AMPK-T184A/57C10; *lkb1^{T2}* (red, $n = 24$), 57C10; *lkb1^{T2}* (yellow, $n = 24$), UAS-AMPK-T184A; *lkb1^{T2}* (blue, $n = 23$) and wt (black, $n = 24$) flies. c) Sleep profiles of UAS-AMPK/57C10; *lkb1^{T2}* (red, $n = 24$), 57C10; *lkb1^{T2}* (yellow, $n = 24$), UAS-AMPK; *lkb1^{T2}* (blue, $n = 11$) and wt (black, $n = 24$) flies. (d–g) Statistical analysis of sleep duration, sleep bout number, sleep bout duration, and latency to sleep in wt (black, $n = 24$), 57C10; *lkb1^{T2}* (yellow, $n = 24$), UAS-AMPK-T184E; *lkb1^{T2}* (blue, $n = 22$), UAS-AMPK-T184E/57C10; *lkb1^{T2}* (red, $n = 19$), UAS-AMPK-T184A; *lkb1^{T2}* (blue, $n = 23$), UAS-AMPK-T184A/57C10; *lkb1^{T2}* (red, $n = 24$), UAS-AMPK; *lkb1^{T2}* (blue, $n = 11$), and UAS-AMPK/57C10; *lkb1^{T2}* (red, $n = 24$) flies. Open bars denote daytime, filled bars nighttime; n.s. not shown. One-way ANOVA was used. n.s. denotes $P > 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Error bars represent SEM.

find that LKB1 regulates sleep need as indicated by reduced NREM δ power density in *Lkb1* knockdown mutant mice.

Sleep is important for animals. Sleep regulation is accomplished through 2 processes: circadian and sleep homeostatic (Borbely 1982; Borbely et al. 2016). The circadian clock regulates the timing of sleep, and homeostatic process regulates the sleep drive. Molecular mechanisms of the circadian clock have been revealed through research in *Drosophila* and other organisms (Hendricks et al. 2000; Shaw et al. 2000; Nitabach and Taghert 2008; Allada and Chung 2010; Mohawk et al. 2012). Although many sleep-related genes have been identified in sleep regulation (Cirelli 2009; Allada et al. 2017; Jan et al. 2020), our understanding of the mechanisms underlying sleep homeostatic regulation remains limited (Allada et al. 2017; Donlea et al. 2017).

Multiple regions in *Drosophila* and mouse brains have been implicated in sleep regulation. In flies, sleep is regulated by several regions including: the ILNv and DN1 clock neurons which are important for circadian control of sleep (Parisky et al. 2008; Shang et al. 2008; Sheeba et al. 2008; Chung et al. 2009; Shang et al. 2013; Kunst et al. 2014). And the mushroom bodies, the dorsal of fan-shaped body, the ellipsoid body, the pars intercerebralis, and glia (Joiner et al. 2006; Foltenyi et al. 2007; Crocker et al. 2010; Donlea et al. 2011; Guo et al. 2011; Seugnet et al. 2011; Liu et al. 2012; Ueno et al. 2012; Yi et al. 2013; Donlea et al. 2014; Park et al. 2014; Chen et al. 2015; Liu et al. 2016; Pimentel et al. 2016). In mammals, sleep is regulated by monoaminergic, cholinergic, glutamatergic, and GABAergic neurons that are distributed in




Fig. 5. Genetic interactions between *lkb1* and *sik3*. a) Sleep profiles of *lkb1*^{T2} (red, n = 45), *Sik3-T196E-flag; lkb1*^{T2} (dark red, n = 46), *Sik3-flag; lkb1*^{T2} (orange, n = 48), *Sik3-T196E-flag* (blue, n = 45), *Sik3-flag* (dark blue, n = 44) and wt (black, n = 46) flies. (b–e) Statistical analysis of sleep duration, sleep bout number, sleep bout duration, and latency to sleep in wt (black, n = 46), *Sik3-flag* (dark blue, n = 44), *Sik3-T196E-flag* (blue, n = 45), *Sik3-flag; lkb1*^{T2} (orange, n = 48), *Sik3-T196E-flag; lkb1*^{T2} (dark red, n = 46), and *lkb1*^{T2} (red, n = 45) flies. Open bars denote daytime, filled bars nighttime. Statistics for groups *Sik3-flag; lkb1*^{T2} and *Sik3-T196E-flag; lkb1*^{T2} were presented. One-way ANOVA was used. n.s. denotes P > 0.05, *P < 0.05, **P < 0.01, ***P < 0.001. Error bars represent SEM.

multiple regions including the brain stem, the preoptic hypothalamus, the lateral hypothalamus, and the basal forebrain (Weber and Dan 2016; Saper and Fuller 2017; Scammell et al. 2017). It will be interesting to investigate whether *Lkb1* functions in all or a limited subset of neurons to regulate sleep.

Lkb1 as a master kinase can regulate the activities of ARKs by phosphorylating the site in the active T loop equivalent to AMPK-T172 (Lizcano et al. 2004). Because both AMPK and SIK3 are involved in sleep regulation, it will be interesting to investigate downstream kinases mediating the function of *Lkb1* in sleep regulation. Is it SIK3, AMPK, or other members of the ARKs? Our findings of allele-specific genetic interactions between *Lkb1* and *Ampk* suggest that they could be upstream and downstream of each other in

regulating sleep. Because of the lethality of double mutation combination of *Sik3* and *lkb1*, we cannot rule out that *Sik3* may also be downstream of *Lkb1* in regulating *Drosophila* sleep. The Ca²⁺/calmodulin-dependent protein kinase kinase-2 (CaMKK2, also known as CaMKKβ) could phosphorylate AMPKα-T172 (Hawley et al. 2005; Hurley et al. 2005; Woods et al. 2005; Anderson et al. 2008), but CaMKK2 could not phosphorylate the equivalent sites in the other ARKs, including SIK3 (Fogarty et al. 2010). It will be interesting to investigate whether and how CaMKK2 regulates sleep.

In *Drosophila*, LKB1 functions through SIK3 which phosphorylates histone deacetylase 4 (HDAC4) to regulate lipid storage (Choi et al. 2015). It will be interesting to investigate whether HDAC4 is downstream of LKB1 in sleep regulation.




Fig. 6. Sleep phenotypes of *lkb1* conditional knockout mice. a) Levels of LKB1 protein from *Lkb1*^{fl/fl} mice injected with AAV-hSyn-Cre-GFP virus (Cre⁺ *Lkb1*^{fl/fl}, *n* = 4), *Lkb1*^{fl/fl} mice injected with AAV-hSyn-GFP virus (GFP⁺ *Lkb1*^{fl/fl}, *n* = 3), *Lkb1*^{fl/+} mice injected with AAV-hSyn-Cre-GFP virus (Cre⁺ *Lkb1*^{fl/+}, *n* = 2) and *Lkb1*^{+/+} mice injected with AAV-hSyn-Cre-GFP virus (Cre⁺ *Lkb1*^{+/+}, *n* = 2). These mice were among those used for EEG recording and analysis. b) Levels of LKB1 protein in individual mice (genotypes labeled: Cre⁺ *Lkb1*^{fl/fl}, GFP⁺ *Lkb1*^{fl/fl}, Cre⁺ *Lkb1*^{fl/+}, and Cre⁺ *Lkb1*^{+/+}). These mice were the same mice as those in (a) but presented individually. Statistical analysis of wake duration, NREM duration and REM duration in Cre⁺ *Lkb1*^{fl/fl} (red, *n* = 10), GFP⁺ *Lkb1*^{fl/fl} (yellow, *n* = 5), Cre⁺ *Lkb1*^{fl/+} (blue, *n* = 7), and Cre⁺ *Lkb1*^{+/+} (black, *n* = 4) mice in a 12:12 LD cycle. White background denotes daytime, gray background nighttime. c) Wake duration. Daytime wake duration of Cre⁺ *Lkb1*^{fl/fl} mice was higher than those of GFP⁺ *Lkb1*^{fl/fl}, Cre⁺ *Lkb1*^{fl/+}, or Cre⁺ *Lkb1*^{+/+} mice. Night-time wake duration of Cre⁺ *Lkb1*^{fl/fl} mice was higher than that of GFP⁺ *Lkb1*^{fl/fl} mice. e) NREM duration. Daytime NREM duration of Cre⁺ *Lkb1*^{fl/fl} mice was lower than those of GFP⁺ *Lkb1*^{fl/fl}, Cre⁺ *Lkb1*^{fl/+}, and Cre⁺ *Lkb1*^{+/+} mice. Night-time NREM duration of Cre⁺ *Lkb1*^{fl/fl} mice was lower than that of GFP⁺ *Lkb1*^{fl/fl} mice. g) REM duration. Daytime and nighttime REM durations of Cre⁺ *Lkb1*^{fl/fl} mice was not significantly different from those of GFP⁺ *Lkb1*^{fl/fl}, Cre⁺ *Lkb1*^{fl/+}, and Cre⁺ *Lkb1*^{+/+} mice. EEG power spectrum of (d) WAKE, (f) NREM, and (h) REM states in Cre⁺ *Lkb1*^{fl/fl} (red, *n* = 8), GFP⁺ *Lkb1*^{fl/fl} (yellow, *n* = 5), Cre⁺ *Lkb1*^{fl/+} (blue, *n* = 6), and Cre⁺ *Lkb1*^{+/+} (black, *n* = 4) mice. (i) NREM δ -power density of Cre⁺ *Lkb1*^{fl/fl} (red, *n* = 8), GFP⁺ *Lkb1*^{fl/fl} (yellow, *n* = 5), Cre⁺ *Lkb1*^{fl/+} (blue, *n* = 6), and Cre⁺ *Lkb1*^{+/+} (black, *n* = 4) mice. One-way ANOVA was used in (c, e, g) for comparison of Cre⁺ *Lkb1*^{fl/fl}, Cre⁺ *Lkb1*^{fl/+}, and Cre⁺ *Lkb1*^{+/+} mice. Unpaired t test was used in (c, e, g) for comparison of Cre⁺ *Lkb1*^{fl/fl}, and GFP⁺ *Lkb1*^{fl/fl} mice. Two-way ANOVA followed by Turkey's multiple comparisons test was used in (d, f, h, i). n.s. denotes *P* > 0.05, **P* < 0.05, ***P* < 0.01, ****P* < 0.001. Error bars represent SEM.

More importantly, an important question for further studies is whether Lkb1 regulation of sleep is related to its regulation of metabolism. Changes in energy homeostasis directly and reversibly influence the sleep/wake cycle (Collet *et al.* 2016). Some molecules involved in metabolism regulate sleep (Taheri *et al.* 2004; Bjorness and Greene 2009; Thimgan *et al.* 2010; Gerstner *et al.* 2011; Nixon *et al.* 2015; Grubbs *et al.* 2020). In *Drosophila*, starvation suppresses sleep without building up sleep drive (Thimgan *et al.* 2010). Lkb1 and its downstream components are involved in regulating metabolism, with examples such as LKB1-AMPK signaling in the liver regulating glucose homeostasis (Shaw *et al.* 2005), SIK3-HDAC4 regulating energy balance in *Drosophila* (Wang *et al.* 2011). Either LKB1 has 2 independent roles in sleep and metabolism or that its 2 roles are related.

Our recent *in vitro* biochemical discoveries of STE20 phosphorylation of AMPK and SIK3 (and other ARKs) raise more questions about physiological significance of any STE20 or any other ARK in sleep (Liu, Wang, Cui, Gao, *et al.* 2022; Liu, Wang, Cui, *et al.* 2022).

Data availability

Strains and plasmids are available upon request. All data necessary for confirming the conclusions of the article are present within the article and its [supplementary data](#).

[Supplemental material](#) is available at GENETICS online.

Acknowledgments

We are grateful to Dr Bowen Deng for Ampk flies, to Ping-ping Yan, Lan Wang, and Yong-hui Zhang for technical assistance, to Drs Wei Yang and En-xin Zhou for *Drosophila* video tracing programs, to Dan Wang for mouse rearing, to members of the Rao lab for discussion, to the Bloomington *Drosophila* Stock Center for flies, to the Jackson Laboratory for mice, to CIBR, Peking-Tsinghua Center for Life Sciences, IDG/McGovern Institute for Brain Research at Peking University and Changping Laboratory for support.

Conflicts of interest

The authors declare no conflict of interests.

Literature cited

- Alessi DR, Sakamoto K, Bayascas JR. LKB1-dependent signaling pathways. *Annu Rev Biochem.* 2006;75:137–163.
- Allada R, Chung BY. Circadian organization of behavior and physiology in *Drosophila*. *Annu Rev Physiol.* 2010;72:605–624.
- Allada R, Cirelli C, Sehgal A. Molecular mechanisms of sleep homeostasis in flies and mammals. *Cold Spring Harb Perspect Biol.* 2017;9(8):a027730.
- Anderson KA, Ribar TJ, Lin F, Noeldner PK, Green MF, Muehlbauer MJ, Witters LA, Kemp BE, Means AR. Hypothalamic CaMKK2 contributes to the regulation of energy balance. *Cell Metab.* 2008;7(5):377–388.
- Bardeesy N, Sinha M, Hezel AF, Signoretti S, Hathaway NA, Sharpless NE, Loda M, Carrasco DR, DePinho RA. Loss of the Lkb1 tumour suppressor provokes intestinal polyposis but resistance to transformation. *Nature.* 2002;419(6903):162–167.
- Barger Z, Frye CG, Liu D, Dan Y, Bouchard KE. Robust, automated sleep scoring by a compact neural network with distributional shift correction. *PLoS One.* 2019;14(12):e0224642.
- Beg ZH, Allmann DW, Gibson DM. Modulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase activity with cAMP and with protein fractions of rat liver cytosol. *Biochem Biophys Res Commun.* 1973;54(4):1362–1369.
- Bjorness TE, Greene RW. Adenosine and sleep. *Curr Neuropharmacol.* 2009;7(3):238–245.
- Borbely AA. A two process model of sleep regulation. *Hum Neurobiol.* 1982;1(3):195–204.
- Borbely AA, Baumann F, Brandeis D, Strauch I, Lehmann D. Sleep deprivation: effect on sleep stages and EEG power density in man. *Electroencephalogr Clin Neurophysiol.* 1981;51(5):483–495.
- Borbely AA, Daan S, Wirz-Justice A, Deboer T. The two-process model of sleep regulation: a reappraisal. *J Sleep Res.* 2016;25(2):131–143.
- Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development.* 1993;118(2):401–415.
- Carling D, Clarke PR, Zammit VA, Hardie DG. Purification and characterization of the AMP-activated protein-kinase. Copurification of acetyl-CoA carboxylase kinase and 3-hydroxy-3-methylglutaryl-CoA reductase kinase-activities. *Eur J Biochem.* 1989;186(1–2):129–136.
- Carling D, Zammit VA, Hardie DG. A common bicyclic protein-kinase cascade inactivates the regulatory enzymes of fatty-acid and cholesterol-biosynthesis. *FEBS Lett.* 1987;223(2):217–222.
- Carlson CA, Kim KH. Regulation of hepatic acetyl coenzyme A carboxylase by phosphorylation and dephosphorylation. *J Biol Chem.* 1973;248(1):378–380.
- Carretero J, Medina PP, Pio R, Montuenga LM, Sanchez-Cespedes M. Novel and natural knockout lung cancer cell lines for the LKB1/STK11 tumor suppressor gene. *Oncogene.* 2004;23(22):4037–4040.
- Chan KY, Jang MJ, Yoo BB, Greenbaum A, Ravi N, Wu WL, Sanchez-Guardado L, Lois C, Mazmanian SK, Deverman BE, *et al.* Engineered AAVs for efficient noninvasive gene delivery to the central and peripheral nervous systems. *Nat Neurosci.* 2017;20(8):1172–1179.
- Chen WF, Maguire S, Sowcik M, Luo W, Koh K, Sehgal A. A neuron-glia interaction involving GABA transaminase contributes to sleep loss in sleepless mutants. *Mol Psychiatry.* 2015;20(2):240–251.
- Chikahisa S, Fujiki N, Kitaoka K, Shimizu N, Sei H. Central AMPK contributes to sleep homeostasis in mice. *Neuropharmacology.* 2009;57(4):369–374.
- Choi S, Lim DS, Chung J. Feeding and fasting signals converge on the LKB1-SIK3 pathway to regulate lipid metabolism in *Drosophila*. *PLoS Genet.* 2015;11(5):e1005263.
- Chu VT, Graf R, Wirtz T, Weber T, Favret J, Li X, Petsch K, Tran NT, Sieweke MH, Berek C, *et al.* Efficient CRISPR-mediated mutagenesis in primary immune cells using CrispRGold and a C57BL/6 Cas9 transgenic mouse line. *Proc Natl Acad Sci U S A.* 2016;113(44):12514–12519.
- Chung BY, Kilman VL, Keath JR, Pitman JL, Allada R. The GABA(a) receptor EDL acts in peptidergic PDF neurons to promote sleep in *Drosophila*. *Curr Biol.* 2009;19(5):386–390.
- Cirelli C. The genetic and molecular regulation of sleep: from fruit flies to humans. *Nat Rev Neurosci.* 2009;10(8):549–560.
- Collet TH, van der Klaauw AA, Henning E, Keogh JM, Suddaby D, Dachi SV, Dunbar S, Kelway S, Dickson SL, Farooqi IS, *et al.* The sleep/wake cycle is directly modulated by changes in energy balance. *Sleep.* 2016;39(9):1691–1700.

- Crocker A, Shahidullah M, Levitan IB, Sehgal A. Identification of a neural circuit that underlies the effects of octopamine on sleep: wake behavior. *Neuron*. 2010;65(5):670–681.
- Daan S, Beersma DG, Borbely AA. Timing of human sleep: recovery process gated by a circadian pacemaker. *Am J Physiol*. 1984;246(2 Pt 2):R161–R183.
- Dai X, Zhou E, Yang W, Mao R, Zhang W, Rao Y. Molecular resolution of a behavioral paradox: sleep and arousal are regulated by distinct acetylcholine receptors in different neuronal types in *Drosophila*. *Sleep*. 2021;44(7):0161–8105.
- Dai X, Zhou E, Yang W, Zhang X, Zhang W, Rao Y. D-serine made by serine racemase in *Drosophila* intestine plays a physiological role in sleep. *Nat Commun*. 2019;10(1):2041–1723.
- Davies SP, Hawley SA, Woods A, Carling D, Haystead TAJ, Hardie DG. Purification of the AMP-activated protein-kinase on ATP-gamma-sepharose and analysis of its subunit structure. *Eur J Biochem*. 1994;223(2):351–357.
- Deng B, Li Q, Liu X, Cao Y, Li B, Qian Y, Xu R, Mao R, Zhou E, Zhang W, et al. Chemoconnectomics: mapping chemical transmission in *Drosophila*. *Neuron*. 2019;101(5):876–893.e4.
- Dijk DJ, Beersma DG, Daan S. EEG power density during nap sleep: reflection of an hourglass measuring the duration of prior wakefulness. *J Biol Rhythms*. 1987;2(3):207–219.
- Donlea JM, Alam MN, Szymusiak R. Neuronal substrates of sleep homeostasis; lessons from flies, rats and mice. *Curr Opin Neurobiol*. 2017;44:228–235.
- Donlea JM, Pimentel D, Miesenbock G. Neuronal machinery of sleep homeostasis in *Drosophila*. *Neuron*. 2014;81(4):860–872.
- Donlea JM, Thimgan MS, Suzuki Y, Gottschalk L, Shaw PJ. Inducing sleep by remote control facilitates memory consolidation in *Drosophila*. *Science*. 2011;332(6037):1571–1576.
- Ferrer A, Caelles C, Massot N, Hegardt FG. Activation of rat-liver cytosolic 3-hydroxy-3-methylglutaryl coenzyme A reductase kinase by adenosine 5'-monophosphate. *Biochem Biophys Res Commun*. 1985;132(2):497–504.
- Fogarty S, Hawley SA, Green KA, Saner N, Mustard KJ, Hardie DG. Calmodulin-dependent protein kinase kinase-beta activates AMPK without forming a stable complex: synergistic effects of Ca²⁺ and AMP. *Biochem J*. 2010;426(1):109–118.
- Foltenyi K, Greenspan RJ, Newport JW. Activation of EGFR and ERK by rhomboid signaling regulates the consolidation and maintenance of sleep in *Drosophila*. *Nat Neurosci*. 2007;10(9):1160–1167.
- Franken P, Chollet D, Tafti M. The homeostatic regulation of sleep need is under genetic control. *J Neurosci*. 2001;21(8):2610–2621.
- Funato H, Miyoshi C, Fujiyama T, Kanda T, Sato M, Wang Z, Ma J, Nakane S, Tomita J, Ikkyu A, et al. Forward-genetics analysis of sleep in randomly mutagenized mice. *Nature*. 2016;539(7629):378–383.
- Gerstner JR, Vanderheyden WM, Shaw PJ, Landry CF, Yin JCP. Fatty acid binding proteins modulate sleep and enhance long-term memory consolidation in *Drosophila*. *PLoS One*. 2011;6(1):e15890.
- Gill RK, Yang SH, Meerzaman D, Mechanic LE, Bowman ED, Jeon HS, Chowdhuri SR, Shakoori A, Dracheva T, Hong KM, et al. Frequent homozygous deletion of the LKB1/STK11 gene in non-small cell lung cancer. *Oncogene*. 2011;30(35):3784–3791.
- Grubbs JJ, Lopes LE, van der Linden AM, Raizen DM. A salt-induced kinase is required for the metabolic regulation of sleep. *PLoS Biol*. 2020;18(4):e3000220.
- Guldberg P, thor Straten P, Ahrenkiel V, Seremet T, Kirkin AF, Zeuthen J. Somatic mutation of the Peutz-Jeghers syndrome gene, LKB1/STK11, in malignant melanoma. *Oncogene*. 1999;18(9):1777–1780.
- Guo F, Yi W, Zhou MM, Guo AK. Go signaling in mushroom bodies regulates sleep in *Drosophila*. *Sleep*. 2011;34(3):273–281.
- Han C, Jan LY, Jan YN. Enhancer-driven membrane markers for analysis of nonautonomous mechanisms reveal neuron-glia interactions in *Drosophila*. *Proc Natl Acad Sci U S A*. 2011;108(23):9673–9678.
- Hardie DG. AMP-activated protein kinase: maintaining energy homeostasis at the cellular and whole-body levels. *Annu Rev Nutr*. 2014;34:31–55.
- Hardie DG, Schaffer BE, Brunet A. AMPK: an energy-sensing pathway with multiple inputs and outputs. *Trends Cell Biol*. 2016;26(3):190–201.
- Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Makela TP, Alessi DR, Hardie DG. Complexes between the LKB1 tumor suppressor, STRAD alpha/beta and MO25 alpha/beta are upstream kinases in the AMP-activated protein kinase cascade. *J Biol*. 2003;2(4):28.
- Hawley SA, Davison M, Woods A, Davies SP, Beri RK, Carling D, Hardie DG. Characterization of the AMP-activated protein kinase kinase from rat liver and identification of threonine 172 as the major site at which it phosphorylates AMP-activated protein kinase. *J Biol Chem*. 1996;271(44):27879–27887.
- Hawley SA, Pan DA, Mustard KJ, Ross L, Bain J, Edelman AM, Frenguelli BG, Hardie DG. Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab*. 2005;2(1):9–19.
- Hearle N, Schumacher V, Menko FH, Olschwang S, Boardman LA, Gille JJ, Keller JJ, Westerman AM, Scott RJ, Lim W, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res*. 2006;12(10):3209–3215.
- Hemminki A, Markie D, Tomlinson I, Avizienyte E, Roth S, Loukola A, Bignell G, Warren W, Aminoff M, Hoglund P, et al. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature*. 1998;391(6663):184–187.
- Hemminki A, Tomlinson I, Markie D, Jarvinen H, Sistonen P, Bjorkqvist AM, Knuutila S, Salovaara R, Bodmer W, Shibata D, et al. Localization of a susceptibility locus for Peutz-Jeghers syndrome to 19p using comparative genomic hybridization and targeted linkage analysis. *Nat Genet*. 1997;15(1):87–90.
- Hendricks JC, Finn SM, Panckeri KA, Chavkin J, Williams JA, Sehgal A, Pack AI. Rest in *Drosophila* is a sleep-like state. *Neuron*. 2000;25(1):129–138.
- Herzig S, Shaw RJ. AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol*. 2018;19(2):121–135.
- Honda T, Fujiyama T, Miyoshi C, Ikkyu A, Hotta-Hirashima N, Kanno S, Mizuno S, Sugiyama F, Takahashi S, Funato H, et al. A single phosphorylation site of SIK3 regulates daily sleep amounts and sleep need in mice. *Proc Natl Acad Sci U S A*. 2018;115(41):10458–10463.
- Hong SP, Leiper FC, Woods A, Carling D, Carlson M. Activation of yeast Snf1 and mammalian AMP-activated protein kinase by upstream kinases. *Proc Natl Acad Sci U S A*. 2003;100(15):8839–8843.
- Hurley RL, Anderson KA, Franzone JM, Kemp BE, Means AR, Witters LA. The Ca²⁺/calmodulin-dependent protein kinase kinases are AMP-activated protein kinase kinases. *J Biol Chem*. 2005;280(32):29060–29066.
- Ingebritsen TS, Lee HS, Parker RA, Gibson DM. Reversible modulation of the activities of both liver microsomal hydroxymethylglutaryl coenzyme A reductase and its inactivating enzyme. Evidence for regulation by phosphorylation-dephosphorylation. *Biochem Biophys Res Commun*. 1978;81(4):1268–1277.
- Jan M, O'Hara BF, Franken P. Recent advances in understanding the genetics of sleep. *F1000Research*. 2020;9:214.

- Jeghers H, Mc KV, Katz KH. Generalized intestinal polyposis and melanin spots of the oral mucosa, lips and digits; a syndrome of diagnostic significance. *N Engl J Med.* 1949;241(26):1031–1036.
- Jenne DE, Reimann H, Nezu J, Friedel W, Loff S, Jeschke R, Muller D, Back W, Zimmer M. Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat Genet.* 1998;18(1):38–44.
- Ji HB, Ramsey MR, Hayes DN, Fan C, McNamara K, Kozlowski P, Torrice C, Wu MC, Shimamura T, Perera SA, et al. LKB1 modulates lung cancer differentiation and metastasis. *Nature.* 2007;448(7155):807–U807.
- Jishage K, Nezu J, Kawase Y, Iwata T, Watanabe M, Miyoshi A, Ose A, Habu K, Kake T, Kamada N, et al. Role of Lkb1, the causative gene of Peutz-Jegher's syndrome, in embryogenesis and polyposis. *Proc Natl Acad Sci U S A.* 2002;99(13):8903–8908.
- Joiner WJ, Crocker A, White BH, Sehgal A. Sleep in *Drosophila* is regulated by adult mushroom bodies. *Nature.* 2006;441(7094):757–760.
- Klarsfeld A, Leloup JC, Rouyer F. Circadian rhythms of locomotor activity in *Drosophila*. *Behav Process.* 2003;64(2):161–175.
- Kunst M, Hughes ME, Raccuglia D, Felix M, Li M, Barnett G, Duah J, Nitabach MN. Calcitonin gene-related peptide neurons mediate sleep-specific circadian output in *Drosophila*. *Curr Biol.* 2014;24(22):2652–2664.
- Liu QL, Liu S, Kodama L, Driscoll MR, Wu MN. Two dopaminergic neurons signal to the dorsal fan-shaped body to promote wakefulness in *Drosophila*. *Curr Biol.* 2012;22(22):2114–2123.
- Liu S, Liu QL, Tabuchi M, Wu MN. Sleep drive is encoded by neural plastic changes in a dedicated circuit. *Cell.* 2016;165(6):1347–1360.
- Liu YX, Wang TV, Cui Y, Gao S, Rao Y. Biochemical purification uncovers mammalian sterile 3 (MST3) as a new protein kinase for multifunctional protein kinases AMPK and SIK3. *J Biol Chem.* 2022;298(5):101929.
- Liu YX, Wang TV, Cui Y, Li C, Jiang L, Rao Y. STE20 phosphorylation of AMPK related kinases revealed by biochemical purifications combined with genetics. *J Biol Chem.* 2022;298(5):101928.
- Lizcano JM, Goransson O, Toth R, Deak M, Morrice NA, Boudeau J, Hawley SA, Udd L, Makela TP, Hardie DG, et al. LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. *EMBO J.* 2004;23(4):833–843.
- Lopez M, Dieguez C. Cellular energy sensors: AMPK and beyond. *Mol Cell Endocrinol.* 2014;397(1–2):1–3.
- Martin SG, St Johnston D. A role for *Drosophila* LKB1 in anterior-posterior axis formation and epithelial polarity. *Nature.* 2003;421(6921):379–384.
- Matsumoto S, Iwakawa R, Takahashi K, Kohno T, Nakanishi Y, Matsuno Y, Suzuki K, Nakamoto M, Shimizu E, Minna JD, et al. Prevalence and specificity of LKB1 genetic alterations in lung cancers. *Oncogene.* 2007;26(40):5911–5918.
- Mehenni H, Gehrig C, Nezu J, Oku A, Shimane M, Rossier C, Guex N, Blouin JL, Scott HS, Antonarakis SE. Loss of LKB1 kinase activity in Peutz-Jeghers syndrome, and evidence for allelic and locus heterogeneity. *Am J Hum Genet.* 1998;63(6):1641–1650.
- Michell BJ, Stapleton D, Mitchelhill KI, House CM, Katsis F, Witters LA, Kemp BE. Isoform-specific purification and substrate specificity of the 5'-AMP-activated protein kinase. *J Biol Chem.* 1996;271(45):28445–28450.
- Mitchelhill KI, Stapleton D, Gao G, House C, Michell B, Katsis F, Witters LA, Kemp BE. Mammalian AMP-activated protein-kinase shares structural and functional homology with the catalytic domain of yeast Snf1 protein-kinase. *J Biol Chem.* 1994;269(4):2361–2364.
- Miyoshi H, Nakau M, Ishikawa T, Seldin MF, Oshima M, Taketo MM. Gastrointestinal hamartomatous polyposis in Lkb1 heterozygous knockout mice. *Cancer Res.* 2002;62(8):2261–2266.
- Mohawk JA, Green CB, Takahashi JS. Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci.* 2012;35:445–462.
- Morton JP, Jamieson NB, Karim SA, Athineos D, Ridgway RA, Nixon C, Mckay CJ, Carter R, Brunton VG, Frame MC, et al. Lkb1 haploinsufficiency cooperates with KRAS to promote pancreatic cancer through suppression of p21-dependent growth arrest. *Gastroenterology.* 2010;139(2):586–597.
- Munday MR, Carling D, Hardie DG. Negative interactions between phosphorylation of acetyl-coA carboxylase by the cyclic AMP-dependent and AMP-activated protein-kinases. *FEBS Lett.* 1988;235(1–2):144–148.
- Nagy S, Maurer GW, Hentze JL, Rose M, Werge TM, Rewitz K. AMPK signaling linked to the schizophrenia-associated 1q21.1 deletion is required for neuronal and sleep maintenance. *PLoS Genet.* 2018;14(12):e1007623.
- Nakada D, Saunders TL, Morrison SJ. Lkb1 regulates cell cycle and energy metabolism in haematopoietic stem cells. *Nature.* 2010;468(7324):653–658.
- Nitabach MN, Taghert PH. Organization of the *Drosophila* circadian control circuit. *Curr Biol.* 2008;18(2):R84–93.
- Nixon JP, Mavanji V, Butterick TA, Billington CJ, Kotz CM, Teske JA. Sleep disorders, obesity, and aging: the role of orexin. *Ageing Res Rev.* 2015;20:63–73.
- Parisky KM, Agosto J, Pulver SR, Shang YH, Kuklin E, Hodge JLL, Kang K, Liu X, Garrity PA, Rosbash M, et al. PDF cells are a GABA-responsive wake-promoting component of the *Drosophila* sleep circuit. *Neuron.* 2008;60(4):672–682.
- Park M, Miyoshi C, Fujiyama T, Kakizaki M, Ikkyu A, Honda T, Choi J, Asano F, Mizuno S, Takahashi S, et al. Loss of the conserved PKA sites of SIK1 and SIK2 increases sleep need. *Sci Rep.* 2020;10(1):8676.
- Park S, Sonn JY, Oh Y, Lim C, Choe J. SIFamide and SIFamide receptor define a novel neuropeptide signaling to promote sleep in *Drosophila*. *Mol Cells.* 2014;37(4):295–301.
- Peutz JLA. Very remarkable case of familial polyposis of mucous membrane of intestinal tract and nasopharynx accompanied by peculiar pigmentations of skin and mucous membrane. *Nederl Maandschr Geneesk.* 1921;10:134–146.
- Pimentel D, Donlea JM, Albot CBT, Ong SMS, Hurston AJFT, Miesenbock G. Operation of a homeostatic sleep switch. *Nature.* 2016;536(7616):333–337.
- Poe AR, Wang B, Sapor ML, Ji H, Li K, Onabajo T, Fazliyeva R, Gibbs M, Qiu Y, Hu Y, et al. Robust CRISPR/Cas9-mediated tissue-specific mutagenesis reveals gene redundancy and perdurance in *Drosophila*. *Genetics.* 2019;211(2):459–472.
- Qian Y, Cao Y, Deng B, Yang G, Li J, Xu R, Zhang D, Huang J, Rao Y. Sleep homeostasis regulated by 5HT2b receptor in a small subset of neurons in the dorsal fan-shaped body of *Drosophila*. *Elife.* 2017;6:e26519.
- Rowan A, Bataille V, MacKie R, Healy E, Bicknell D, Bodmer W, Tomlinson I. Somatic mutations in the Peutz-Jeghers (LKB1/STK11) gene in sporadic malignant melanomas. *J Invest Dermatol.* 1999;112(4):509–511.
- Sakamoto K, McCarthy A, Smith D, Green KA, Hardie DG, Ashworth A, Alessi DR. Deficiency of LKB1 in skeletal muscle prevents AMPK activation and glucose uptake during contraction. *EMBO J.* 2005;24(10):1810–1820.
- Sanchez-Cespedes M, Parrella P, Esteller M, Nomoto S, Trink B, Engles JM, Westra WH, Herman JG, Sidransky D. Inactivation of

- LKB1/STK11 is a common event in adenocarcinomas of the lung. *Cancer Res.* 2002;62(13):3659–3662.
- Saper CB, Fuller PM. Wake-sleep circuitry: an overview. *Curr Opin Neurobiol.* 2017;44:186–192.
- Sato F, Muragaki Y, Zhang YP. DEC1 negatively regulates AMPK activity via LKB1. *Biochem Biophys Res Commun.* 2015;467(4):711–716.
- Scammell TE, Arrigoni E, Lipton JO. Neural circuitry of wakefulness and sleep. *Neuron.* 2017;93(4):747–765.
- Sengupta S, Nagalingam A, Muniraj N, Bonner MY, Mistriotis P, Afthinos A, Kuppusamy P, Lanoue D, Cho S, Korangath P, et al. Activation of tumor suppressor LKB1 by honokiol abrogates cancer stem-like phenotype in breast cancer via inhibition of oncogenic Stat3. *Oncogene.* 2017;36(41):5709–5721.
- Seugnet L, Suzuki Y, Merlin G, Gottschalk L, Duntley SP, Shaw PJ. Notch signaling modulates sleep homeostasis and learning after sleep deprivation in *Drosophila*. *Curr Biol.* 2011;21(10):835–840.
- Shackelford DB, Shaw RJ. The lkb1-ampk pathway: metabolism and growth control in tumour suppression. *Nat Rev Cancer.* 2009;9(8):563–575.
- Shang YH, Donelson NC, Vecsey CG, Guo F, Rosbash M, Griffith LC. Short neuropeptide f is a sleep-promoting inhibitory modulator. *Neuron.* 2013;80(1):171–183.
- Shang YH, Griffith LC, Rosbash M. Light-arousal and circadian photoreception circuits intersect at the large PDF cells of the *Drosophila* brain. *Proc Natl Acad Sci U S A.* 2008;105(50):19587–19594.
- Shaw PJ, Cirelli C, Greenspan RJ, Tononi G. Correlates of sleep and waking in *Drosophila melanogaster*. *Science.* 2000;287(5459):1834–1837.
- Shaw RJ, Kosmatka M, Bardeesy N, Hurley RL, Witters LA, DePinho RA, Cantley LC. The tumor suppressor LKB1 kinase directly activates amp-activated kinase and regulates apoptosis in response to energy stress. *Proc Natl Acad Sci U S A.* 2004;101(10):3329–3335.
- Shaw RJ, Lamia KA, Vasquez D, Koo SH, Bardeesy N, DePinho RA, Montminy M, Cantley LC. The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science.* 2005;310(5754):1642–1646.
- Sheeba V, Fogle KJ, Kaneko M, Rashid S, Chou YT, Sharma VK, Holmes TC. Large ventral lateral neurons modulate arousal and sleep in *Drosophila*. *Curr Biol.* 2008;18(20):1537–1545.
- Shen Z, Wen XF, Lan F, Shen ZZ, Shao ZM. The tumor suppressor gene LKB1 is associated with prognosis in human breast carcinoma. *Clin Cancer Res.* 2002;8(7):2085–2090.
- Skoulidis F, Byers LA, Diao LX, Papadimitrakopoulou VA, Tong P, Izzo J, Behrens C, Kadara H, Parra ER, Canales JR, et al. Co-occurring genomic alterations define major subsets of KRAS-mutant lung adenocarcinoma with distinct biology, immune profiles, and therapeutic vulnerabilities. *Cancer Discov.* 2015;5(8):860–877.
- Sutherland CM, Hawley SA, McCartney RR, Leech A, Stark MJR, Schmidt MC, Hardie DG. Elm1p is one of three upstream kinases for the *Saccharomyces cerevisiae* SNF1 complex. *Curr Biol.* 2003;13(15):1299–1305.
- Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index (BMI). *Sleep.* 2004;27:146–147.
- Tanwar PS, Mohapatra G, Chiang S, Engler DA, Zhang LH, Kaneko-Tarui T, Ohguchi Y, Birrer MJ, Teixeira JM. Loss of LKB1 and PTEN tumor suppressor genes in the ovarian surface epithelium induces papillary serous ovarian cancer. *Carcinogenesis.* 2014;35(3):546–553.
- Thimgan MS, Suzuki Y, Seugnet L, Gottschalk L, Shaw PJ. The perilipin homologue, lipid storage droplet 2, regulates sleep homeostasis and prevents learning impairments following sleep loss. *PLoS Biol.* 2010;8(8):e1000466.
- Tobler I, Borbely AA. Sleep EEG in the rat as a function of prior waking. *Electroencephalogr Clin Neuro.* 1986;64(1):74–76.
- Tomlinson IP, Houlston RS. Peutz-Jeghers syndrome. *J Med Genet.* 1997;34(12):1007–1011.
- Ueno T, Tomita J, Tanimoto H, Endo K, Ito K, Kume S, Kume K. Identification of a dopamine pathway that regulates sleep and arousal in *Drosophila*. *Nat Neurosci.* 2012;15(11):1516–1523.
- Wang B, Moya N, Niessen S, Hoover H, Mihaylova MM, Shaw RJ, Yates JR, III, Fischer WH, Thomas JB, Montminy M. A hormone-dependent module regulating energy balance. *Cell.* 2011;145(4):596–606.
- Weber F, Dan Y. Circuit-based interrogation of sleep control. *Nature.* 2016;538(7623):51–59.
- Werth E, Dijk DJ, Achermann P, Borbely AA. Dynamics of the sleep EEG after an early evening nap: experimental data and simulations. *Am J Physiol.* 1996;271(3 Pt 2):R501–510.
- Westerman AM, Entius MM, de Baar E, Boor PP, Koole R, van Velthuysen ML, Offerhaus GJ, Lindhout D, de Rooij FW, Wilson JH. Peutz-Jeghers syndrome: 78-year follow-up of the original family. *Lancet.* 1999;353(9160):1211–1215.
- Wingo SN, Gallardo TD, Akbay EA, Liang MC, Contreras CM, Boren T, Shimamura T, Miller DS, Sharpless NE, Bardeesy N, et al. Somatic LKB1 mutations promote cervical cancer progression. *PLoS One.* 2009;4(4):e5137.
- Woods A, Dickerson K, Heath R, Hong SP, Momcilovic M, Johnstone SR, Carlson M, Carling D. Ca²⁺/calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab.* 2005;2(1):21–33.
- Woods A, Johnstone SR, Dickerson K, Leiper FC, Fryer LG, Neumann D, Schlattner U, Wallimann T, Carlson M, Carling D. LKB1 is the upstream kinase in the AMP-activated protein kinase cascade. *Curr Biol.* 2003;13(22):2004–2008.
- Yardeni T, Eckhaus M, Morris HD, Huizing M, Hoogstraten-Miller S. Retro-orbital injections in mice. *Lab Anim (NY).* 2011;40(5):155–160.
- Yeh LA, Kim KH. Regulation of acetyl-coA carboxylase - properties of coA activation of acetyl-coA carboxylase. *Proc Natl Acad Sci U S A.* 1980;77(6):3351–3355.
- Yi W, Zhang YP, Tian YJ, Guo J, Li Y, Guo AK. A subset of cholinergic mushroom body neurons requires go signaling to regulate sleep in *Drosophila*. *Sleep.* 2013;36(12):1809–1821.
- Yurgel ME, Kakad P, Zandawala M, Nassel DR, Godenschwege TA, Keene AC. A single pair of leucokinin neurons are modulated by feeding state and regulate sleep-metabolism interactions. *PLoS Biol.* 2019;17(2):e2006409.
- Zhang X, Yan H, Luo Y, Huang Z, Rao Y. Thermoregulation-independent regulation of sleep by serotonin revealed in mice defective in serotonin synthesis. *Mol Pharmacol.* 2018;93(6):657–664.

Communicating editor: H. Bellen